

Where to next in limited-stage small cell lung cancer?

● ● ● ● ● ● ● ● ● ●
The role of immune checkpoint inhibitors

Disclaimer

- *Unapproved products or unapproved uses of approved products may be discussed by the faculty; these situations may reflect the approval status in one or more jurisdictions*
- *The presenting faculty have been advised by USF Health and touchIME to ensure that they disclose any such references made to unlabelled or unapproved use*
- *No endorsement by USF Health and touchIME of any unapproved products or unapproved uses is either made or implied by mention of these products or uses in USF Health and touchIME activities*
- *USF Health and touchIME accept no responsibility for errors or omissions*

Expert MDT faculty



Dr Lizza Hendriks

Maastricht University Medical Center
Maastricht, The Netherlands



Dr Jacob Sands

Dana-Farber Cancer Institute
Boston, MA, USA



Prof. Suresh Senan

Amsterdam University Medical Center
Amsterdam, The Netherlands

Experience of using ICIs in SCLC

ES-SCLC

1L PD-1/PD-L1 inhibitors + ChT

CAPSTONE-1 (N=462) adebrelimab + ChT¹
Significantly improved mOS vs placebo + ChT
(p=0.0008)

IMpower133 (N=403) atezolizumab + ChT²
Significantly improved mOS vs placebo + ChT
(HR 0.76; 95% CI 0.60–0.95; p=0.0154).
Atezolizumab + ChT 5-year OS estimate of 12%
(n=18)³

CASPIAN (N=805) durvalumab + ChT⁴
Significantly improved mOS vs ChT alone
(p=0.0003)

ANSTRUM-005 (N=585) serplulimab + ChT⁵
Significantly improved mOS vs placebo + ChT
(p<0.001)

LS-SCLC

STIMULI trial⁶

Consolidation nivolumab + ipilimumab vs
observation after CRT in LS-SCLC (N=222)

mPFS not significant: 10.7 vs 14.5 months
(HR 1.02; p=0.93)
Median follow-up: 22.4 months

mOS not significant: NR vs 32.1 months
(HR 0.95; p=0.82)
Median follow-up: 35 months

**STIMULI trial did not meet its primary
endpoint of improvement in PFS with
nivolumab–ipilimumab consolidation
after CRT**

Pembrolizumab + concurrent CRT⁷

Single institution study of pembrolizumab +
concurrent CRT (N=40)

Adverse events:
grade 5 events, n=0; grade 4 events, n=3;
grade 3 events, n=41

Median follow-up: 23.1 months

mPFS: 19.7 months


mOS: 39.5 months

**Concurrent CRT and pembrolizumab
was well tolerated and OS and PFS
were favourable compared to those in
the CONVERT trial**

1L, first-line; ChT, chemotherapy; CI, confidence interval; CRT, chemoradiotherapy; ES-SCLC, extensive-stage SCLC; HR, hazard ratio; ICI, immune checkpoint inhibitors; LS-SCLC, limited-stage SCLC; m, median; NR, not reached; OS, overall survival; PD-1, programmed cell death protein 1; PD-L1, programmed cell death ligand 1; PFS, progression free survival; SCLC, small cell lung cancer. 1. Cheng Y, et al. *Immuno-oncol Technol.* 2023;20:100556; 2. Liu SV, et al. *J Clin Oncol.* 2021;39:619–30; 3. Reck M, et al. *Lung Cancer.* 2024;196:107924; 4. Paz-Ares L, et al. *ESMO Open.* 2022;7:100408; 5. Cheng Y, et al. *JAMA.* 2022;328:1223–32; 6. Peters S, et al. *Ann Oncol.* 2022;33:67–79; 7. Welsh JW, et al. *J Thorac Oncol.* 2020;15:1919–27.

Ongoing trials of ICIs in LS-SCLC

Phase III trials



Phase II trials

- **Atezolizumab** (ACHILES; NCT03540420)
- **Durvalumab** (DURVALUNG; NCT05617963)
- **Envafohimab** (NCT05904015)
- **Tislelizumab** (NeoSCLC-001; NCT06375109)
- **TQB2450** (NCT04539977)

Given in combination with CRT

Consolidation therapy
(When disease has not progressed following CRT)

Phase III efficacy and safety data

Adebrelimab (NCT04691063)¹

Adebrelimab with cCRT followed by addebrelimab maintenance (data cut-off: 31 Oct 2023)*

Primary endpoint: Safety (N=28)

	% of patients
Grade ≥3 TRAEs	96.4% [†]
TR pneumonitis	14.3% (all grade 2)
TR immune-mediated lung disease	3.6% (grade 2)

Efficacy

ORR	92.9%
mPFS	17.9 months
mOS	NR

Adebrelimab with cCRT showed acceptable tolerability and favourable efficacy outcomes

Durvalumab (ADRIATIC)²

Durvalumab ± tremelimumab vs placebo consolidation (data cut-off: 15 Jan 2024)

Primary endpoint: Efficacy

	D (n=264)	P (n=266)	P
mOS [‡] (months)	55.9	33.4	p=0.01
mPFS [§] (months)	16.6	9.2	p=0.02

Safety

	D (n=262)	P (n=265)
Grade 3/4 AEs	24.4%	24.2%
Pneumonitis	38.2%	30.2%
Grade 3/4 pneumonitis	3.1%	2.6%

Durvalumab consolidation treatment was well tolerated and demonstrated improvements in OS and PFS vs placebo

Atezolizumab (NRG-LU005)³

CRT + atezolizumab followed by atezolizumab maintenance vs CRT alone[¶]

Primary endpoint: Efficacy

	A + CRT (n=274)	CRT (n=270)	P
mOS (months)	33.1	39.5	p=0.7640
mPFS (months)	12.0	11.5	p=0.9542

Safety^{**}

	A + CRT (n=267)	CRT (n=254)
Grade 3/4 AEs	86.5%	92.5%
Grade 3/4 irAEs	15.7%	6.2%
Pneumonitis	26.2%	11.8%

Concurrent atezolizumab did not improve survival vs standard CRT

Direct comparisons between trials should not be made due to differences in trial design. *Median follow-up 29.4 months; †all events with incidence of ≥10% were haematological toxicities;

‡median duration of follow-up: 37.2 months; §median duration of follow-up 27.4 months; ||refers to pneumonitis and radiation pneumonitis collectively; ¶median follow-up 21.0 months;

**reporting window 30 days post CRT and 90 days post end of atezolizumab (11 weeks vs 15 months). A, atezolizumab; AE, adverse event; cCRT, concurrent CRT; CRT, chemoradiotherapy;

D, durvalumab; irAE, immune-related AE; m, median; NR, not reached; ORR, objective response rate; OS, overall survival; P, placebo; PFS, progression free survival; TR, treatment-related.

1. Cheng Y et al. Presented at: ELCC Congress 2024, Prague, Czech Republic. 20–23 March 2024. Abstr. 198P; 2. Cheng Y, et al. *N Engl J Med.* 2024; 391:1313–27;

3. Higgins KA, et al. Presented at: 2024 ASTRO Annual Meeting, Washington, DC, USA. 20 September–2 October 2024. Abstr. LBA02.

Phase II efficacy and safety data

Serplulimab (ASTRUM-LC01)¹

Single arm: serplulimab consolidation therapy after cCRT and PCI (data cut-off: 7 April 2024)*

Primary endpoint: Efficacy (N=55)

ORR	96.4%
DCR	96.4%
mPFS	NR

Safety (% of patients)

TRAE	72.7%
Grade ≥3 TRAEs	14.6%
Most common grade 3/4 AE	Pneumonitis

Serplulimab consolidation therapy demonstrated response rates of >95% and a manageable safety profile

Toripalimab (GASTO-1052A; NCT04418648)²

Toripalimab consolidation vs observation with CR or PR after cCRT[†]

Primary endpoint: Efficacy

	T (n=31)	O (n=33)	
2-year PFS	61.6%	34.8%	p=0.04
2-year OS	82.7%	59.1%	p=0.23

Safety (% of patients)

	T (n=31)	O (n=33)
Grade ≥2 pneumonitis	16.1%	9.1%
Grade ≥4 TRAEs	0	0

Toripalimab consolidation following definitive cCRT was effective and tolerable in LS-SCLC

Camrelizumab (ChiCTR2000032275)³

Camrelizumab plus ChT followed by cCRT plus camrelizumab vs CRT[‡]

Primary endpoint: Efficacy

	Camrelizumab (n=17)	CRT (n=17)
1-year PFS	54.5%	44.4%
mPFS (months)	NR	14.4

Safety (% of patients)

	Camrelizumab (n=17)	CRT (n=17)
Grade 3/4 TRAEs	58.8%	52.9%
Grade 3/4 radiation pneumonitis	5.9%	5.9%

Addition of camrelizumab to CRT resulted in an encouraging 1-year PFS rate and did not increase incidence of AEs

Direct comparisons between trials should not be made due to differences in trial design. *Median follow-up 9.8 months; †median follow-up 25 months; ‡median follow-up 10.6 months. AE, adverse event; ChT, chemotherapy; cCRT, concurrent CRT; CR, complete response; CRT, chemoradiotherapy; DCR, disease control rate; LS-SCLC, limited-stage small cell lung cancer; m, median; NR, not reached; O, observation; ORR, objective response rate; OS, overall survival; PCI, prophylactic cranial irradiation; PFS, progression free survival; PR, partial response; T, toripalimab; TRAE, treatment-related AE. 1. Wu Y et al. Presented at: ESMO Congress 2024, Barcelona, Spain. 13–17 September 2024. Poster 1759P; 2. Zhang PX, et al. Presented at: ASCO Annual Meeting 2024, Chicago, IL, USA. 31 May–4 June 2024. Abstr. P8098; 3. Hu M, et al. Presented at: WCLC 2024, San Diego, CA, USA. 7–10 September 2024. Poster P1.13A.05.